

Report for 2003ND25B: Comparative Study of Fossil and Extant Fish Growth: Including Analyses of Mean Annual Temperature in the Geologic Record

- Conference Proceedings:
 - Newbrey, M.G. and M.A. Bozek. 2002. Growth of *Esox tiemani* of the Paleocene Sentinel Butte Formation of North Dakota with evidence to estimate mean annual temperature. Society of Vertebrate Paleontology 62nd Annual Meeting, Vol. 22.
 - Newbrey, M.G. and A.C. Ashworth. 2003. Fish community dynamics, growth of yellow perch, and correlations with climate and fire in an early Holocene lake in North Dakota. Society of Vertebrate Paleontology 63rd Annual Meeting, Vol. 23.
 - Poster: Newbrey, M.G. and A.C. Ashworth. July 2003. A new method to estimate early Holocene mean annual temperature using growth characteristics of fossil *Perca flavescens* (yellow perch). XVI International Union for Quaternary Research Congress.
- Other Publications:
 - Presentation: Newbrey, Michael and Allan Ashworth, October 2003, FISH COMMUNITY DYNAMICS, GROWTH OF YELLOW PERCH, AND CORRELATIONS WITH CLIMATE AND FIRE IN AN EARLY HOLOCENE LAKE IN NORTH DAKOTA, Society of Vertebrate Paleontology Meetings, St. Paul, MN
 - Presentation: Michael Newbrey, Sept. 23, 2003, Comparative Study of Fossil and Extant Fish Growth: Including Analyses of Mean Annual Temperature in the Geologic Record, ND WRII Advisory Committee Meeting, Bismarck, North Dakota.
- Articles in Refereed Scientific Journals:
 - Manuscript accepted for publication: Newbrey, M.G. and A.C. Ashworth. A fossil record of colonization and response of lacustrine fish populations to climate change. Canadian Journal of Fisheries and Aquatic Sciences.

Report Follows

COMPARATIVE STUDY OF FOSSIL AND EXTANT FISH GROWTH: INCLUDING ANALYSES OF MEAN ANNUAL TEMPERATURE IN THE GEOLOGIC RECORD

ND WRI Graduate Research Fellowship Project
Michael Newbrey, Fellow
Allan Ashworth, Advisor and Principal Investigator
Department of Geological Sciences
North Dakota State University
Fargo, ND 58105

DESCRIPTION OF THE REGIONAL WATER PROBLEM

It is important to consider the implications of climatic change on surface water resources in light of potential consequences of global warming. North Dakota boasts some of the best long-term data sets in the form of a fossil record to measure the effect of climatic warming on a single population of fish. A fossil lake bed near Jamestown, ND will provide perhaps thousands of years of continuous data of fish growth during a warming climate. This research will provide insight for fishery biologists and wetland ecologists concerning the long-term response of contemporary fish growth in North Dakota given potential climatic changes.

LITERATURE SUMMARY AND PRIOR WORK

Population dynamics have rarely been studied from ancient ecosystems. However, fossils can provide information about growth, mortality, and numerous other ecological processes that have been examined in extant systems (Ricker, 1975). A recent study which detailed the ageing of *Joffrichthys triangulperus* (Newbrey and Bozek, 2000) scales and the subsequent modeling of their growth provided insight into the population dynamics of an extinct osteoglossid fish (Newbrey and Bozek, In Press). However, there are no contemporary species within the genus *Joffrichthys* to use as a reference for a comparative growth analysis. More recently, I have conducted research with an extinct form of pike (*Esox tiemani*) (Wilson, 1980) and compared its growth to living forms of *Esox* (i.e., northern pike and muskellunge). The research was presented at the 2002 Society of Vertebrate Paleontology meetings and will be submitted to the journal *Palaeogeography, Palaeoclimatology, Palaeoecology* for publication. The manuscript details a correlation between growth and mean annual temperature and introduces a new procedure to estimate mean annual temperature in the fossil record. Much of the methodology presented in this proposal is derived from my research of pike growth.

Climatological processes strongly influence growth rates of fish. For example, growth can vary depending on aquatic and ambient mean annual temperatures (Gillooly et al. 2002). Air temperature strongly influences surface water temperature (McCombie, 1959; Livingstone and Lotter, 1998) and therefore fish growth. Temperature has a strong influence on metabolic rate (Q_{10} relationship) and growth on ectothermic taxa; a temperature increase of 10°C increases the metabolic rate two to three fold and reaction rates increase 100-200% (Cossins and Bowler, 1987). Our pike research indicates that temperature is correlated with latitude. The pike grow more slowly in the northerly latitudes because of cooler and shorter growing seasons and more quickly in the warmer-longer growing seasons of southern latitudes.

The Fellowship research will focus on a comparative growth of several groups of fish. Specifically, we will contrast growth of fish in the fossil record to that of living fish to determine mean annual temperature change in the geologic record. The analyses are important for fishery biologists and ecologists in North Dakota who are interested in the implications of climatic change on surface water

resources and fish. For example, we will study a fossil glacial lake site in North Dakota that has produced fossil specimens of contemporary species of fish. The environment of the fossil lake changed from a cool wet climate with tamarack, black spruce, birch and aspen to a contemporary prairie-pothole region. The change occurred over a period of thousands of years thus giving us insight into ecological processes that are affected by current climate changes.

A site of particular interest in North Dakota represents a glacial, 10,000 year old fossil lake bed containing extremely well preserved fish (yellow perch, *Perca flavescens*), which are still living in ND today. Some of the best preserved Late Wisconsinan/Early Holocene fossils ever reported, come from a 10,000 year-old site (Seibold site) underlying an ephemeral wetland on the Missouri Coteau, near Buchanan, North Dakota. Cvancara et al. (1971) described the geology and paleolimnology of one lake deposit called the Seibold site (SW $\frac{1}{4}$ NW $\frac{1}{4}$ Sec. 21, T141N, R67W), Stutsman County, North Dakota. The Seibold site represents the remains of a small basin (100m in diameter), which was probably part of large lake complex during the Late Wisconsinan/Early Holocene. The site has excellent preservation of terrestrial and aquatic plants and animals, which permits taxonomic assignments of the fossil remains to the generic and specific levels. Study of plant specimens have revealed that the lake was initially surrounded by a spruce-dominated assemblage that was replaced by a mixed assemblage of deciduous trees, shrub, herbs, and grasses (Cvancara et al., 1971). Animals recovered from the site include five species of fish, frogs, muskrat, and evidence of beaver (Cvancara et al., 1971)(see fossils <http://www.ndsu.nodak.edu/instruct/ashworth/SEIBOLD.pdf>).

The changes in pollen indicate this site is ideal for a long-term study of fish growth in a North Dakota lake, where all data can be gathered in one excavation. During the period of lake sediment deposition, while the climate was warming and the glacier was retreating to Canada, the botanical environment around the lake changed (Cvancara et al., 1971). Fossil pollen indicates that the plant community changed from a forest to prairie ecosystem around 7,500 years ago. As the lake warmed over a few thousand years, measurable change in patterns of growth should be detectable for yellow perch. Numerous studies have modeled growth of individual populations of yellow perch (i.e., Scott and Crossman, 1973; Becker, 1983; etc.) and, in all publications combined, from a variety of latitudes and ambient mean annual temperatures providing a spectrum of growth curves to assemble and analyze. The combined growth curves will provide a correlation between growth and mean annual temperature and thus provide an index to estimate mean annual temperature change in the fossil record. This study site and others in this research will provide insight to fishery biologists and surface water ecologists about the effects of temperature change on fish populations in North Dakota and other areas.

SCOPE AND OBJECTIVES

The overall objective of this Fellowship research is to describe changes in the paleoclimate and fossil fish growth by comparing fossil specimens to extant fish populations. For example, some site-specific objectives include: 1.) to conduct an excavation at the Seibold site near Jamestown, ND to collect fossil yellow perch; 2.) to describe growth of fossil yellow perch; 3.) to compile and analyze contemporary yellow perch growth data in order to contrast growth in relation to the fossils; and 4.) to analyze contemporary yellow perch growth in relation to ambient mean annual temperature in order to calculate a chronological series of estimates of ambient mean annual temperature.

These objectives will be repeated for numerous other localities, represented in museum collections, and containing the following taxa: bowfin, *Amia calva*; the mooneyes and goldeyes of *Hiodon*; and the members of the pikes, *Esox*. These genera are represented in the present North American fish assemblage, in the fossil record from the Cretaceous to the present, and in museum collections.

METHODS, PROCEDURES, AND FACILITIES

Numerous publications contain the information needed to locate specimens (e.g., Grande, 1984; Grande and Bemis, 1998; Grande, 1999; Li et al., 1997; Wilson, 1980, 1981, 1984; etc.) suitable for data. Typically, hard structures (i.e., scales, otoliths, cleithra, fin rays, spines, and vertebrae) from captured fish are aged and growth is calculated using the relation between growth of these structures and fish total length (Van Oosten, 1941; Carlander, 1969). Total length is the distance from the anterior-most tip of the head to the vertical plane of the posterior caudal fin tips. For this study, fossil scales and vertebrae will be aged in the lab and from museum collections by counting annulus marks or light and dark pairs of bands on bones. Fishery biologists use Von Bertalanffy growth curves (Von Bertalanffy, 1938) to assess growth rates. The curve is fit to the maximum total lengths for each age class:

$$TL_t = L_\infty [1 - e^{-K(t-t_0)}]$$

where:

TL_t = Total length (cm) at t (age in years);

L_∞ = maximum total length;

K = the Brody growth coefficient;

t = time (i.e., age in years);

t_0 = time at age zero (time at theoretical zero length).

Yellow perch growth data will be taken from Scott and Crossman (1973), Becker (1983), and a list of other sources to contrast extant growth curves. Because male and female yellow perch have different growth rates and life spans all data from extant populations will be combined for both sexes. This is done to standardize error to match that of the fossil material. Site specific mean annual temperature (MAT) data rounded to the nearest 0.1°C will be taken from the WorldClimate© web site and when possible checked for accuracy with Northern Oceanic and Atmospheric Administration (NOAA) data. Ambient MAT will be used in a linear regression analysis and regressed to the natural log transformed total lengths (cm) of age three of yellow perch to obtain a correlation between MAT and growth:

$$\text{Ln}(TL_{\text{Age}3}) = m \times \sqrt{\text{MAT}_E + 10^\circ\text{C}} + b$$

where:

$\text{Ln}(TL_{\text{Age}3})$ = natural log transformed total length (TL cm) at 3 years of age;

m = slope parameter of linear regression;

MAT_E = mean annual temperature (MAT) at sites of extant (E) populations;

b = intercept parameter of linear regression.

Von Bertalanffy total length will be used because it includes the growth parameters and describes more variation in growth than individual equation parameters alone. Age three total lengths will be used for analysis because this age represents a stable portion of the curve where the effects of the parameters L_∞ and t_0 are reduced while still providing a range of sizes to contrast prior to asymptotic length. To estimate the ambient MAT of the fossil site, the linear regression equation was algebraically rearranged:

$$\text{MAT}_F = \left(\frac{\text{Ln}(TL_{\text{Age}3}) - b}{m} \right)^2 - 10^\circ\text{C}$$

where:

MAT_F = mean annual temperature (MAT) of the fossil (F) site;

$\text{Ln}(TL_{\text{Age}3})$ = natural log transformed total length (TL cm) at age three;

b = intercept parameter of linear regression;
 m = slope parameter of linear regression.

ANTICIPATED RESULTS

As the glacial lake matured to a modern pothole wetland and as the climate changed from a cool moist environment to a dry warmer environment there were quantifiable changes occurring in fish growth. Because there is a correlation between ambient air temperature and water temperature within the first meter of depth (McCombie 1959; Livingstone and Lotter 1998), we hypothesize that growth of fossil yellow perch will initially be slow in the cool climate and then increase as the climate warmed. Mean annual temperature change will be modeled from contemporary fish growth by a mathematical relation to temperature. The correlation will then be used to predict mean annual temperature change over a few thousand years. This information will provide a better understanding of evolution of natural processes to biologists studying the pothole region in North Dakota. New climatic information will be of interest to researchers studying changes in temperature and wet/dry cycles. The information and analyses obtained from the glacial lake will be presented at a scientific meeting and will ultimately be submitted to a peer-reviewed journal for publication. The procedures will be repeated for the other taxa of fish being examined in my dissertation in order to better understand relations between changes in MAT and fish growth.

RESULTS AND CONCLUSIONS

We show 1) drought was occurring during the late Pleistocene when other studies have only shown this to be a relatively stable / wet period, 2) during the late Pleistocene and early Holocene fish abundance was negatively responding to drought, 3) fish abundance was described by nutrient levels that also respond to the intensity of drought, and 4) the pattern of post-glacial fish colonization of the Missouri Coteau in ND can be better understood from the description of a warming climate and the life-history characteristics of the fishes. This research provides peer-reviewed analyses and interpretation that will be used in a proposal to the National Science Foundation for further research at the site.

I have been aging yellow perch from the Seibold site to estimate more post-glacial paleotemperatures. However, most of the specimens in the UND collection have degraded or lack substrate depth data to be useful for a fish growth in relation to climate change analysis. A growth analysis will only take place if further funding can be procured for an excavation. I have also identified three additional fish from the site since the original research report in 1970: *Stizostedion*, *Catostomus*, and *Phoxinus neogaesius*. All three taxa are still found in North Dakota today.

This research also addresses the trends in fish movement across latitude and through time. I have constructed the largest fossil fish database to date, which is nearly double the size of previous fish reviews published in 1981 and 1986 and consists of nearly 400 taxonomic entries. Preliminary results are very interesting and suggest that fish dispersal is tracking with climate.

I have completed the datasets and analyses for seven of eight of my 'extant' dissertation taxa. These analyses include the growth of five taxa significantly responding to mean annual temperature. Two taxa do not have enough published data available to analyze. Three of the four species of pikes and yellow perch and goldeye growth significantly respond to temperature. Manuscripts are in progress describing the results at a variety of degrees of completion.

REFERENCES

- Becker, G. C. 1983. Fishes of Wisconsin. University of Wisconsin Press, Madison, Wisconsin. 1052 pp.
- Carlander, K. D. 1969. Handbook of freshwater fishery biology, volume one. Iowa State University Press, Ames, Iowa. 752 pp.
- Cossins, A. H. and K. Bowler. 1987. Temperature Biology of Animals. Chapman and Hall, London. 339 pp.
- Cvancara, A. M., Clayton, L., Bickley, W. B., Jacob, A. F., Ashworth, A. C., Brophy, J. A., Shay, C. T., Delorme, L. D., and Lammers, G. E. 1971. Paleolimnology of Late Quaternary deposits: Seibold site, North Dakota. Science 171:269-306.
- Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M., and J. H. Brown. 2002. Effects of size and temperature on developmental time. Nature 417:70-73.
- Grande, L. 1984. Paleontology of the Green River Formation, with a review of the fish fauna. The Geologic Survey of Wyoming, Bulletin 3:1-334. (2nd ed.)
- and W. E. Bemis. 1998. A comprehensive phylogenetic study of amiid fishes (Amiidae) based on comparative skeletal anatomy. An empirical search for interconnected patterns of natural history. Society of Vertebrate Paleontology Memoir 4:i-x, 1-690; supplement to Journal of Vertebrate Paleontology 18(1).
- King, J. R., B. J. Shuter, and A. P. Zimmerman. 1999. Empirical links between thermal habitat, fish growth, and climate change. Transactions of the American Fisheries Society 128:656-665.
- Li G.-q., M. V. H. Wilson, and L. Grande 1997. Review of *Eohiodon* (Teleostei: Osteoglossomorpha) from western North America, with a phylogenetic reassessment of Hiodontidae. Journal of Paleontology 71:1109-1124.
- Livingstone, D. M. and A. F. Lotter. 1998. The relationship between air and water temperatures in lakes of the Swiss Plateau: a case study with palaeolimnological implications. Journal of Paleolimnology 19:181-198.
- McCombie, A. M. 1959. Some relations between air temperatures and the surface water temperatures of lakes. Limnology and Oceanography 4:252-258.
- Newbrey, M. G., and M. A. Bozek. 2000. A new species of *Joffrichthys* (Teleostei: Osteoglossidae) from the Sentinel Butte Formation, (Paleocene) of North Dakota, USA. Journal of Vertebrate Paleontology 20:12-20.
- Newbrey, M. G., and M. A. Bozek. In Press. Age, growth, and mortality of *Joffrichthys triangulpterus* (Teleostei: Osteoglossidae) from the Paleocene Sentinel Butte Formation, North Dakota, USA. Journal of Vertebrate Paleontology 00:00-00.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. Bulletin of the Fisheries Research Board of Canada Bulletin 191:1-382.
- Van Oosten, J. 1941. The age and growth of freshwater fishes; pp. 196-205 in A symposium on hydrobiology. University of Wisconsin Press, Madison, Wisconsin. 405 pp.
- Scott, W. B. and E. J. Crossman. 1973. Freshwater fishes of Canada. Research Board of Canada Bulletin 184.
- Von Bertalanffy, L. 1938. A quantitative theory of organic growth. Human Biology 10:181-213.

- Wilson, M. V. H. 1980. Oldest known *Esox* (Pices: Esocidae), part of a new Paleocene teleost fauna from western Canada. *Canadian Journal of Earth Sciences* 17:307-312.
- 1981. Eocene Freshwater fishes from the Coalmont Formation, Colorado. *Journal of Paleontology* 55:671-674.
- 1984. Year classes and sexual dimorphism in the Eocene catostomid fish *Amyzon aggregatum*. *Journal of Vertebrate Paleontology* 3:137-142.